

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

questionable validity in the light of Osterhout's researches in this line.<sup>32</sup>—C. R. B.

Anatomy of the ovule of Myrica.—Miss Kershaw<sup>33</sup> has investigated the ovule of Myrica Gale, and has discovered that in all of the morphological features it is an ordinary angiosperm, with its solitary megaspore mother cell, linear tetrad, eight-nucleate embryo sac, and porogamy. The following anatomical features, however, are worthy of mention: the nucellus is not only completely free from the single integument but is also distinctly stalked within it; vascular strands (eight or nine in number) traverse the integument, without branching, almost to the apex of the ovule. These two features of the ovule are usually regarded as primitive, belonging to the ancient gymnosperms rather than to angiosperms.—J. M. C.

Phototropism of roots.—LINSBAUER and VOUK, after overcoming many experimental difficulties, have found<sup>3+</sup> that the roots of *Raphanus sativus* and *Sinapis alba*, which have been credited with being only negatively phototropic, react positively or negatively according to the intensity of the light. Roots of the former in moist air turn toward light of about 8 candles, while in water they are much less sensitive, no very certain curvatures being obtained until the light was increased to 400 c.p. Sinapis in water, on the contrary, gave the best positive response at 0.2 c.p., and decided negative curvatures at 0.64 c.p. These results support the Müller-Olimanns theory of phototropism.—C. R. B.

Dispersal of seeds by ants.—Weiss<sup>35</sup> has concluded that the gorse (Ulex) and the broom (Sarothamnus) should be included among myrmecochorous plants, along with Chelidonium, Viola, etc. He finds that the seeds have a brightly colored caruncle containing oily food material and resembling in structure and contents the elaiosomes (of Sernander) of other myrmecochorous plants; that ants are particularly attracted by the oil-containing caruncle, and can and will carry about the seeds of gorse; and that the rectilinear distribution of gorse bushes along actual or disused paths or roadways is only paralleled by the distribution of such plants as the celandine along ant-runs.—J. M. C.

Anatomy of Gleichenia.—BOODLE and HILEY<sup>36</sup> have investigated the vascular structure of Gleichenia, a genus interesting on account of its protostelic species.

<sup>32</sup> Bot. Gazette 46:53-55. 1908.

<sup>&</sup>lt;sup>33</sup> KERSHAW, EDITH MAY, The structure and development of the ovule of *Myrica Gale*. Annals of Botany 23:353-362. pl. 24. 1909.

<sup>&</sup>lt;sup>34</sup> LINSBAUER, K., AND VOUK, V., Zur Kenntnis des Heliotropismus der Wurzeln. Ber. Deutsch. Bot. Gesells. **27:**151–156. 1909.

<sup>35</sup> Weiss, F. E., The dispersal of the seeds of the gorse and the broom by ants. New Phytol. 8:81-89. 1909.

<sup>&</sup>lt;sup>36</sup> BOODLE, L. A., AND HILEY, W. E., On the vascular structure of some species of Gleichenia. Annals of Botany 23:419-432. pl. 29. 1909.